



Gene and protein expression in the myometrium in pregnancy and labor.

Michelle Breuiller-Fouche, Guy Germain

► To cite this version:

Michelle Breuiller-Fouche, Guy Germain. Gene and protein expression in the myometrium in pregnancy and labor.. *Reproduction*, 2006, 131 (5), pp.837-50. 10.1530/rep.1.00725 . inserm-00112636

HAL Id: inserm-00112636

<https://www.hal.inserm.fr/inserm-00112636>

Submitted on 4 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Gene and protein expression in myometrium in pregnancy and labor**

2

3 Michelle Breuiller-Fouche¹, Guy Germain^{2,3}

4

5 ¹ Institut National de la Santé et de la Recherche Médicale, Unité 427, Paris, F-75014, France

6 ² Institut National de la Recherche Agronomique, Biologie du Développement et

7 Reproduction, Centre de Recherches de Jouy, Jouy-en-Josas, F-78352, France

8 ³ author for correspondence, guy.germain@jouy.inra.fr, tel. +33134652501, fax

9 +33134652361

Abstract

Microarray technologies widen our comprehension of the major structural and metabolic transformations which affect the myometrium from the very beginning of pregnancy until parturition. The results are coherent with the mass of information which was accumulated previously, primarily on the basis of studies of selected critical factors. They highlight the activation of precise signaling pathways, some of which may have been previously under evaluated. The remodelling and maturation processes that the myometrium undergoes in pregnancy appear clearly as phenomenon's which last during the full course of gestation. Comparatively, the onset of labor is perhaps the phenomenon which remains the least well described by these methods of analysis. Nevertheless, genomic studies constitute a necessary first step of orientation and help establishing new links between the generic signaling pathways that are activated during the normal or pathological gestation. These studies also represent an indicative step that will have to be paralleled, in the future, with the results of the systematic proteomic analysis of the myometrium.

Introduction

Parturition encompasses composite physiological processes that require synchronization of uterine contractions, cervical dilatation and fetal membrane rupture. The mechanisms by which parturition is triggered remain unknown but they are, as all developmental processes, the results of timely coordinated biochemical and physiological steps. Over the past decades, reductionist approaches have investigated the cellular and molecular bases of these developmental changes and focused on assessing the regulation of selected critical factors involved in the onset of labor.

Recently, the near-entire genomic sequence of the human and several model animals have provided the opportunity to enhance our understanding of the relationships between genes, phenotypes and global transcriptional status. Oligonucleotides or DNA microarray technologies allow the examination of the function of thousands of genes at once and in parallel, thereby providing an “assay” of the transcriptional status of cells or tissues in a wide variety of physiological or pathophysiological situations (Dunckley *et al.*, 2005). In the context of uncomplicated or complicated pregnancy and parturition, their interest is to obtain a molecular snapshot of the expression profile of gene transcripts as a function of the time-dependent process regulating myometrial activity.

Functional genomic studies during pregnancy

Functional genomics has arrived at the study of parturition in different species: mice (Bethin *et al.*, 2003, Salomonis *et al.*, 2005), rats (Girotti & Zingg, 2003), human (Bethin *et al.*, 2003, Chan *et al.*, 2002, Charpigny *et al.*, 2003, Esplin *et al.*, 2005, Havelock *et al.*, 2005, Keelan *et*

al., 2003, Marvin *et al.*, 2002a, 2002b, Tashima *et al.*, 2002), and different tissues:
 myometrium (Aguan *et al.*, 2000, Bethin *et al.*, 2003, Chan *et al.*, 2002, Charpigny *et al.*,
 2003, Esplin *et al.*, 2005, Havelock *et al.*, 2005, Salomonis *et al.*, 2005), fetal membranes
 (Chen *et al.*, 2002, Keelan *et al.*, 2003, Marvin *et al.*, 2002a, 2002b, Muhle *et al.*, 2001,
 Tashima *et al.*, 2002), trophoblast (Chen *et al.*, 2002, Kato *et al.*, 2002). Various
 methodologies have been used: Suppression Subtractive Hybridization (SSH) (Chan *et al.*,
 2002) or microarrays for most of the previous cited studies. All investigators have reported
 genes differentially regulated during parturition (Romero *et al.*, 2002) and multiple novel
 candidate markers for preterm labor (Chan *et al.*, 2002, Marvin *et al.*, 2002a, 2002b, Wu *et*
al., 1999) or for premature rupture of membranes (Tashima *et al.*, 2002). They contributed to
 shed light on specific pathophysiological issues like the patterns of expression of cytokines in
 the fetal membranes and decidua particularly in the presence of intrauterine infection (Keelan
et al., 2003) or on the induction of enzymes for prostaglandin synthesis (Bethin *et al.*, 2003).
 Groups of genes with yet unknown functional connection to these pathologies were also
 found coordinately expressed. This was emphasized in animal studies which allow to explore
 the global pattern of gene expression and their co-regulation on the basis of their genomic
 location over the full time-course of myometrial transformation (i.e. pregnancy) or following
 experimentally-controlled infection (Girotti & Zingg, 2003, Muhle *et al.*, 2001, Salomonis *et*
al., 2005, Wu *et al.*, 1999).

Computational methods have significantly helped to the interpretation of gene profiling
 experiments by delineating clusters of genes sharing coherent expression features (Claverie,
 1999). During the last five years, statistical methods and data analysis for array studies have
 progressed enormously and today an “ideal” study should now incorporate i) careful research
 designs (differential expression or cross sectional studies), ii) statistical methods

incorporating background adjustment and normalization of data, assessment of differential expression after determining sample size to control the proportion of positive calls that are false positives (False Discovery rate), iii) hierarchical clustering (e.g. HOPACH method) (Ganesh *et al.*, 2004, Salomonis *et al.*, 2005), iv) functional organization of genes into pathways networks with the aid of integrated databases like the Gene Ontology consortium (Lewis, 2005) or the Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa, 1997, Kanehisa & Goto, 2000) and finally v) integrating genotype, transcription and clinical trait data, e.g. see (Salomonis *et al.*, 2005).

The biases adopted in this review are several. We focused attention on genes expressed essentially in the “normal” human myometrium. One study has reported a direct comparison of the non-pregnant myometrium (NP) with the pregnant human myometrium at term not in labor (TNIL) (Rehman *et al.*, 2003). It gives an overall picture of the changes that the uterine muscle undergoes in its adaptation to the gestation. Two studies have described the changes in gene expression in the myometrium in preterm patients not in labor (PT) versus patients at term not in labor (TNIL) (Bethin *et al.*, 2003, Charpigny *et al.*, 2003) and two additional studies have described the changes in gene expression in the myometrium in patients at term not in labor (TNIL) versus patients at term in labor (TIL) (Esplin *et al.*, 2005, Havelock *et al.*, 2005). Other studies have not been compiled directly, either because they did not provide sufficient precise references on the genes being studied (i.e. lack of GenBank or Unigene references), they considered other components of the utero-fetal unit (i.e. membranes, decidua, cervix, etc.) or they were carried out on animal species. To translate lists of tens or hundreds of genes found to be differentially regulated in the conditions under study into a clearer understanding of the biological phenomena involved, we used both combinations of searches through the literature referenced in public databases and the Onto-Tools software

developed by the Draghici's group at Wayne State University, Detroit (MI), (<http://vortex.cs.wayne.edu/Projects.html>). The Onto-Express module helps recognizing functional profiles (using Gene Ontology terms) for the categories: biochemical function, biological process, cellular role, cellular component, molecular function, and chromosome location (Draghici *et al.*, 2003). The Pathway-Express module helps data mining, by proposing on the basis of a computational method a hierarchical list of several KEGG pathways (Kanehisa, 1997, Kanehisa & Goto, 2000) most likely adjusted to changes observed in microarray experiments (Khatri *et al.*, 2005). Drawbacks and limitations inherent to the use of these tools or of their cognate alternatives for ontological analysis, have been reviewed recently (Khatri & Draghici, 2005).

The variations of myometrium gene expression in the course of gestation could be dispatched in 14 main KEGG pathways (<http://www.genome.jp/kegg/pathway.html>) that were the most representatives for the changes seen in non-pregnant (NP) versus term quiescent pregnant uterus (TNIL), quiescent preterm (PT) versus term quiescent uterus (TNIL) and term quiescent uterus (TNIL) versus contracting uterus at labor (TIL). Tables 1-4 and Figures 1-2 summarize the compiled data which are commented below in more details. Wherever possible, OMIM nomenclature has been adopted in the text and tables (<http://www.ncbi.nlm.nih.gov/omim/>).

Changes in structural and contractile genes during gestation

Actin cytoskeleton, focal adhesion, adherens and tight junctions related genes represent a large subset of genes that are over-expressed in the pregnant human myometrium as compared to the non-pregnant state (Table 1).

Regulation of actin cytoskeleton

Growth and cytoskeletal remodeling of myometrial cells during pregnancy are critical for myometrial functions including those expressed during labor and delivery. At term, signals that initiate labor, ultimately promote a switch in the phenotype in the quiescent uterus to a smooth muscle which becomes spontaneously active, excitable, highly responsive to uterine agonists, and exhibits a high degree of cell-cell coupling. Myometrial cells are rich in actin microfilaments, intermediate filaments and microtubules that allow cells to adapt to a variety of shapes and to carry out coordinated and directed movements. In rat, myometrial expression of alpha-actin is high throughout pregnancy. An increased expression of gamma-actin and its translocation to the membrane is observed in uterine myocytes at late gestation. Therefore, the alteration in myometrial composition of contractile proteins is important to prepare the myometrium for the development of contractions during labor (Shynlova *et al.*, 2005).

RhoA, a member of the Ras superfamily, and its downstream mediator RhoA kinase (ROCK1) are necessary for agonist-induced stress fiber formation (Gogarten *et al.*, 2001). The higher expression of protein kinase C (PKC) isoforms observed in pregnant *versus* non-pregnant myometrium (Rehman *et al.*, 2003) may promote or drive the formation of stress fibers. Additionally, up-regulation of ROCK1 has been demonstrated in human term myometrium (Moore *et al.*, 2000, Rehman *et al.*, 2003). This leads to the idea that increased endogenous ROCK1 activity, resulting in enhanced RhoA-mediated calcium sensitization, is involved in the increased contractility at the time of labor. It is of note that EGF increases the presence of actin and myometrial EGF receptors transcripts are increased at the time of parturition (Charpigny *et al.*, 2003, Gargiulo *et al.*, 1997, Rehman *et al.*, 2003).

Focal adhesion

The transmission of force between the contractile apparatus of the cell and extracellular matrix (ECM) occurs at membrane-associated dense plaques or “focal adhesions”. Focal adhesions consist of clusters of integrins that mediate interactions between the extra- and intracellular environments. The cytoplasmic regions of integrins connect with cytoskeletal elements and signaling components such as Focal Adhesion Kinase (FAK), while the extracellular regions connect to specific extracellular matrix molecules such as collagen, laminin or fibronectin.

Fetal growth imposes mechanical tension on the myometrium at term, which in turn induces activation of FAK leading to focal adhesion turnover and supporting myometrial cell hypertrophy. During late pregnancy, a fall in tyrosine phosphorylation of FAK and stabilization of focal adhesions occur for provoking the cessation of myometrial hypertrophy. Because actin polymerization and the dynamic remodeling of the actin cytoskeleton play key roles in the regulation of myometrial contraction, there is growing evidence that stretching induce labor, probably through a change in the expression of contraction-associated-proteins (CAPs) (Macphee & Lye, 2000). In human term pregnant myometrium, changes in cytoskeletal organization support a role for FAK and other focal adhesion-associated proteins as regulators of actin dynamics. ITGA5 with its known partner ITGB1, is up-regulated in rat myometrium during late pregnancy and labor (Williams *et al.*, 2005) as well as ITGAV in the TNIL human myometrium (Rehman *et al.*, 2003). Therefore they may interact with actin-binding proteins (e.g. actinin and filamin) to form mechanical links to the cytoskeleton. A strong expression of vinculin is concentrated at actin-vinculin focal adhesion sites in human

myometrial cells (Yu & Lopez Bernal, 1998). At the time of parturition, expression of integrins declines and the focal adhesion and related pathways do not change in studies which compare the TNIL and TIL states of pregnancy (see Table 1).

Parturition as an inflammatory process

Cytokine-chemokines interaction

There is a massive influx in the myometrium of neutrophils, macrophages and T-lymphocytes concomitant with the onset of labor (Osman *et al.*, 2003). An increase in cytokines, interleukin-1 (IL-1), IL-6, IL-8 and tumor necrosis factor (TNF)-alpha within tissues of the laboring uterus and cervix is demonstrated.

Chemokines enhance inflammation by inducing chemotaxis and cell activation of inflammatory cells. CXC-Chemokines, attract neutrophils but not macrophages, while CC-Chemokines preferentially induce migration of macrophages. Chemokine transcripts increase in myometrium at term, including CCL13 (MCP-4 or Monocyte chemotactic protein-4), CCL19, CCL21, CXCR4 (NPYRL or neuropeptide Y receptor-like) and CXCR5 (Burkitt lymphoma receptor) (Bethin *et al.*, 2003, Charpigny *et al.*, 2003, Rehman *et al.*, 2003). At the time of labor, a selective increase in CXCL10 (Interferon-inducible protein-10), CCL8 and CC13 is noted (Esplin *et al.*, 2005) (see Table 2). IL-8, MCP-1 and RANTES are regulated by local growth factors and cytokines such as TNF-alpha, interferon-gamma, and IL-1. IL-8 also potentiates the effect of IL-1-induced myometrial contractions through PGE2 production at the time of parturition and up-regulates myometrial TGF-beta receptors expression in vitro, suggesting an additional autocrine-signaling pathway. Therefore, it is clear that coordination

of chemokine-chemokine receptor interactions plays an important role in successful pregnancy (Kayisli *et al.*, 2002).

The TNF receptor super family member Fas and its cognate ligand (FasL) play a role in cell leiomyoma apoptosis (Wang Y. *et al.*, 2002). Therefore increased expression of TNFRSF6 (i.e. Fas receptor) in term myometrium (Table 2) await consideration in the control of the uterine growth process. Elevated expression of TNFRSF11B is observed in human myometrium with or without labor (Esplin *et al.*, 2005, Rehman *et al.*, 2003). A peak in uterine osteoprotegerin (i.e OPG or TNFRSF11B), a soluble membrane of the TNF-alpha receptor family that acts as a negative regulator for receptor activator of nuclear factor-kB (RANK), has been reported during labor in the rat (Girotti & Zingg, 2003).

The gp130 protein is a subunit component of several cytokines receptors including those for LIF. Cytokines sharing the gp130 (or IL6ST) subunits are referred to as IL6 type family of cytokines and signal through JAK/STAT pathway (Heinrich *et al.*, 1998). They play a role in the regulation of gene activation, proliferation and differentiation. Down-regulation of IL6ST and LIF receptor at term (Table 3) may be an indication of the decrease of proliferation processes before the onset of labor.

IFN-gamma inhibits the proliferation of vascular smooth muscle cells and the synthesis of collagens by myofibroblasts. In human myometrial cells, interferon gamma antagonizes IL-1beta-induced prostaglandin-endoperoxide synthase 2 (COX-2 or PTGS2) expression and PGE2 production (Hertelendy *et al.*, 2002). Therefore, the decrease in IFNGR1 transcript observed during the transition from PT to TNIL states of the uterus (Table 2), could be interpreted as an *ad hoc* evolution for labor onset.

225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248

TLR signaling pathway

Toll-like receptors (TLRs) are evolutionarily conserved pathogen-associated microbial patterns and play important roles in innate immunity in mammals. Ten TLRs are evidenced in the uterus but TLR2 and TLR4 mRNA are expressed at the highest levels. Among the TLR-related gene transcripts, SOCS, CD14 and IRAK mRNAs (see Tables 2 and 4) are widely expressed in the myometrium (Nishimura & Naito, 2005). TLR4 mediates induction of PTL in mice treated with lipopolysaccharide (LPS) (Wang H. & Hirsch, 2003). In monocytes, LPS-induced signaling through TLRs, lead to the recruitment of docking proteins such as IL-1 receptor-associated kinase (IRAK) and TNF-alpha receptor-associated factor (TRAF6) leading to the activation of IkappaB kinase (IKK) complex. These pathways in turn activate transcription factor such as NF-kB that coordinates the induction of genes encoding inflammatory mediators (Guha & Mackman, 2001). In human pregnant myometrial cells a positive immunoreactivity for TLR4 is observed and cells exposure to LPS lead to nuclear translocation of the p65 subunit of NFkB (Dallot *et al.*, 2005). The increased expression of CD14 together with increased expression of the catalytic subunit of PI3K in pregnant myometrium may serve to increase host protection against microbial invasion (Rehman *et al.*, 2003). At the end of pregnancy, because some of the TLR-related genes such as IRAK, IKK, MKK have decreased expression compared to earlier stage of the gestation (Charpigny *et al.*, 2003), it is possible that adaptative responses to immune or growth processes take place.

TGF-beta signaling pathway

TGF-beta regulates its own expression, the expression of ECM, of matrix metalloproteinases (MMP), and of tissue inhibitor of matrix metalloproteinases (TIMP) as well as the growth of leiomyomas and of normal myometrial cells (Chegini *et al.*, 1999). TGF-beta receptors mediate their action through multiple pathways, including smad proteins that convey TGF-beta receptor signals from the cell surface to the nucleus, resulting in transcriptional activation of TGF-beta-responsive genes. TGF-beta also activates mitogen-activated protein kinase (MAPK), PKC, and calcium/calmodulin complex, inducing Smad-independent transcriptional responses. One mechanism by which TGF-beta is activated involves thrombospondin (THBS1). THBS1 promotes replication of smooth cells and modifies responses to contractile agents. THBS1 expression and protein in human myometrial tissues are increased during labor and after the administration of OT (Morimoto *et al.*, 1998). Myometrial transcripts of THBS1 and THBS2 increases during pregnancy (Charpigny *et al.*, 2003, Rehman *et al.*, 2003) and peak at the time of labor concomitantly with the expression of TGF-beta (Esplin *et al.*, 2005). It is possible that the control of the TGF-beta pathway in the near term myometrium facilitates the transition of the quiescent uterus towards a contractile state, as well as it contributes to control the uterine growth.

Wnt signalling pathway

Leiomyomas have high levels of both transcripts of WNT5B and of secreted frizzled related protein 1 (SFRP1), a modulator of Wnt signaling (Mangioni *et al.*, 2005). Strong SFRP1 expression under high estrogenic conditions seems to contribute to the development of uterine leiomyomas through the antiapoptotic effect of SFRP1 (Fukuhara *et al.*, 2002). The increased expression of WNT5B in the pregnant myometrium suggests that the Wnt pathway is important in myometrial adaptation to pregnancy by decreasing apoptotic myometrial cell

death (Rehman *et al.*, 2003). Inversely, other genes associated with the inhibition of cell proliferation such as p53 are increased at the end of gestation (Charpigny *et al.*, 2003).

Kinases located at the crossroad of uterine contractility and myometrial cell proliferation

MAPK-signaling pathway

MAPK signaling cascades regulate diverse processes ranging from contraction, proliferation, differentiation, and development. Five families of MAPKs have been defined in mammalian cells: i) extracellular signal-regulated kinases (ERK1 and ERK2), ii) Jun N-terminal kinases (JNK), iii) p38 kinase isozymes. ERK1 and ERK2 are activated by mitogenic stimuli such as growth factors, cytokines and phorbol esters. Members of the JNK family play crucial roles in regulating responses to various stresses and apoptosis. Among the targets of p38 MAPKs are several transcription factors, including NF- κ B, p53 and ATF2, which modulate the expression of genes encoding inflammatory cytokines (Qi & Elion, 2005). MAPKs are involved in inhibiting gap-junction-mediated cellular communication in rat myometrium (Loch-Caruso *et al.*, 2003). Mechanical stretch of the rat uterus stimulates myometrial cell hypertrophy (Douglas *et al.*, 1988) via a mechanism involving integrin/focal adhesion/MAPK cascades (Macphee & Lye, 2000). Activation of MAPKs is necessary for optimal stretch-induced c-fos expression (Oldenhof *et al.*, 2002). In addition, the spatial expression of MAPK p38 and ERK-1/2 in conjunction with ATF2 isoforms within the uterine corpus during pregnancy and labor is likely to be important in preparation of the uterus for labor (Otun *et al.*, 2005).

In the myometrium, the MAPK pathways are also implicated in the induction of PTGS2 expression (Bartlett *et al.*, 1999, Sooranna *et al.*, 2004) and in mediating the effects of prostaglandins (PGs) (Ohmichi *et al.*, 1997), OT (Zhong *et al.*, 2003), endothelin-1 (ET-1) (Kimura *et al.*, 1999) and corticotropin-releasing hormone (CRH) (Grammatopoulos *et al.*, 2000, Papadopoulou *et al.*, 2004). Increasing maternal plasma levels of CRH during the last weeks of pregnancy and the substantial expression of CRH receptors in choriodecidua, placenta and myometrium suggests that this stress hormone plays a role in the control of human parturition (Sehringer *et al.*, 2004).

MAPK activity increases in the rat myometrium from day 15 to day 20 of gestation and declines sharply just before parturition (Robin *et al.*, 2004). During the same interval, a shift in intracellular distribution of the Ras protein precedes the down-regulation of membrane-dependent mitogenic signaling and uterine hypertrophy as pregnancy approaches parturition (Ruzycky, 1998). During RU-486-induced PTL, as previously described for spontaneous labor, ERK phosphorylation levels increase, as does phosphorylation of caldesmon and of 20-kDa MLC. When rats are chronically treated with an agent that prevents ERK activation, the onset of PTL is delayed (Li *et al.*, 2004, Liu *et al.*, 2004).

OT-mediated ERK1/2 activation in myometrial cells involves a phospholipase C (PLC)-independent pathway (Zhong *et al.*, 2003). A cross-talk between growth factors receptors and the estrogen receptor alpha (ESR1) has been proposed. The estrogen response involves estradiol (E2)-ESR1-mediated responses as well as responses resulting from convergence of growth factor and ESR1-initiated activities (Hewitt *et al.*, 2005). Functional signaling proximal to IGF-1R is maintained in the ER alpha-KO mouse uterus. ER alpha is necessary

for IGF-1 induction of uterine nuclear proliferative responses, and a cross-talk between IGF-1R and ER signaling pathways exists *in vivo* (Klotz *et al.*, 2002).

Protein kinase C

PKC constitute a multigene family located at the crossroad of two essential uterine functions, namely contractility and cell proliferation. Six isoforms of PKC, the conventional PKC isoforms (alpha, betaI, betaII, and gamma), the novel PKC isoforms (delta, epsilon, theta, eta, lambda/iota) and the atypical PKC isoform (zeta) are evidenced in the human pregnant myometrium. Whereas PRKCA is required for proliferation of human myometrial cells (Eude *et al.*, 2002), only activation of PRKCZ results in actin reorganization and elicits contractions of the human myometrium at the end of pregnancy (Di Liberto *et al.*, 2003).

A balance between T helper (Th1) (pro-inflammatory) / Th2 (anti-inflammatory) cytokine production, has been described at the time of parturition, and a link between TNF-alpha/IL1-beta and premature human childbirth has been proposed (Arntzen *et al.*, 1998). Because, atypical PKC's are important components of the TNF/IL1B signaling pathway that controls NF-κB activation, the implication of the PRKCZ in the control of the onset of labor in women is not surprising. (Ozaki *et al.*, 2003) also demonstrated that the levels of mRNA of PRKCB1 isoform in pregnant human myometrium were greater than those in nonpregnant myometrium, a feature confirmed in genomic studies (Table 3) (Rehman *et al.*, 2003). The levels of the phosphorylated substrate for PKC, CPI-17 which is considered to inhibit myosin light chain phosphatase were also greater in the pregnant myometrium. These results suggest that the PKC-mediated contractile mechanism is augmented in human

myometrium after gestation, and that this augmentation may be attributable to the increased activity of the PRKCB1 isoform and CPI-17.

Phosphatidyl inositol signaling pathway

The PI3-kinase cascade is activated through the binding of the regulatory subunit to phosphorylated tyrosine residues, leading to enhanced activity of the catalytic subunit. PI3 kinase phosphorylates phosphatidylinositol-4,5-bisphosphate (PIP2) to generate PIP3 which interacts with protein kinase Akt or protein kinase B (PKB). Activation of Akt influences many cellular functions through PIP3 binding and phosphorylation by phosphoinositide-dependent kinase 1 (PDK1). It includes cytoskeletal organization, cell growth, motility, proliferation and survival. PDK1 is known to phosphorylate PKC-zeta in the activation loop. PI3 kinase is also activated by direct binding of the catalytic subunit to activated Ras and PI3 kinase cascade activation can lead to the activation of the ERK cascade (Stein & Waterfield, 2000).

Increases in $[Ca^{2+}]_i$ are controlled by multiple signal pathways in myometrium. GPCR-mediated stimulation of the Gq/11 subfamily and subsequent activation of PLC subfamily results in generation of IP3, which triggers release of Ca^{2+} from intracellular stores. Five PLC isoforms: beta 1, beta 2, beta 3, gamma 1, and gamma 2 are detected in human myometrium. OT activates human myometrium by interacting with at least two G proteins and possibly three PLC beta isoforms (Phaneuf *et al.*, 1996). Indeed, the amount of PLCB4 is increased at midpregnancy, whereas PLCB1, PLCB2, and PLCB3 are up-regulated at term in the rat uterus (Mhaouty-Kodja *et al.*, 2004). PLCB3 may be targeted by both contractant and relaxant signaling pathways in myometrium and play a critical role in the balance between

them (Zhong *et al.*, 2005). Both DAG-sensitive PKC, activated by PLCB products, and DAG-insensitive PKC, possibly activated by PI3kinase-dependent process are involved in ERK activation to modulate rat uterine functions (Robin *et al.*, 2002). In addition, protein tyrosine kinase/phosphatase activities may control both phosphorylation and activation of PLCG1 and contribute to the modulation of the generation of inositol phosphates and uterine tension.

Genomic studies exploring the transition from the NP to the TNIL state found an up-regulation of the transcripts pertaining to the phosphoinositol signaling pathway (Rehman *et al.*, 2003), whereas those comparing the PT to the TNIL state found a down-regulation of transcript expression (Bethin *et al.*, 2003, Charpigny *et al.*, 2003). This may be interpreted as a slowdown of myometrial cell proliferation processes when pregnancy comes close to term (Table 3).

Apoptosis and parturition

Proliferation of smooth muscle cells and fibroblasts occurs in early pregnancy and decreases as the uterus nears parturition. In late pregnancy and during involution of the rat uterus, an increase of apoptosis is observed (Leppert, 1998). However, the extent of apoptosis or dedifferentiation that may occur in human myometrial cells during post-partum involution is still unknown. The most investigated factors linked to apoptosis are the Fas ligand (FasL), Fas receptor (FasR), the TNFalpha and its receptors. Some of the genes associated with the inhibition of cell proliferation and differentiation are increased in the myometrium at the end of pregnancy (Table 4). For an example, the TNFR6 or Fas antigen and the p53 tumor suppressor limit cellular proliferation by inducing cell cycle arrest and apoptosis. The presence of DFFA (DNA fragmentation factor-45), known to be essential for chromatin

condensation during normal apoptosis and the inhibition of the PI3kinase in the term pregnant myometrium, may also contribute to an increase of apoptotic events.

G-protein signaling and parturition

The silencing of the myometrial contractile function is conditioned by a predominant functional cAMP/cGMP system, whereas contractions are under the control of agonist-induced calcium mobilization via the PLC pathway. Gestational-related modifications of GPCR in the myometrium, as well as changes in their associated kinases (GRK), cognate G proteins, and effectors have been detected (Hertelendy & Zakar, 2004, Lopez Bernal & TambyRaja, 2000). Some GPCRs coupled to PLC generates the second messengers IP3 and DAG. The ensuing rise in IP3 releases Ca^{2+} from the sarcoplasmic reticulum, causing a sudden rise in intracellular calcium, whereas DAG activates PKC and the MAPK cascades. Uterine contractility can also be enhanced via the activation of Rho GTPases, and the subsequent action of ROCK's to potentiate the effect of myosin light chain kinase (MYLK). These pathways have been discussed above. Other GPCR coupled to the adenylyl cyclase promotes the accumulation of cAMP which controls the relaxation of the myometrium during pregnancy via the inactivation of MYLK and the activation of ATF-2 (Bailey & Europe-Finner, 2005). Another powerful way for increasing cAMP concentration consists in inhibiting its hydrolysis by phosphodiesterases (PDEs). Among the five PDE families (PDE 1, 2, 3, 4 and 5) identified in the human myometrium (Leroy *et al.*, 1994, Leroy *et al.*, 1999), one particular isoform, PDE4B2, which specifically hydrolyses cAMP is selectively induced at the end of pregnancy suggesting a role for this protein in the setting of the contractile state of the myometrium just before delivery (Mehats *et al.*, 2001, Mehats *et al.*, 2000).

Neuroactive-ligands and parturition

A small selection of peptides, neuroactive factors and ion channels such as ET-1, noradrenaline, NPY, PGE2, GABA... that mediate uterine relaxation or contraction and act through GPCR are mentioned here (see Table 4). ET-1 exerts a stimulatory effect on myometrium contractility and proliferation. Its transcript is up-regulated in the pregnant myometrium (Rehman *et al.*, 2003). Among the three subtypes of alpha2-adrenoceptors found in the human myometrium at term pregnancy (Adolfsson *et al.*, 1998), only the alpha2C-subtype appears to be predominant in pregnant myometrium. Stimulation of GABA (A) receptors tonically inhibits contractions of the rabbit myometrium, while stimulation of GABA (B) receptors enhances contractions. Steroids interact with GABA (A) receptors to modulate uterine contractility (Majewska & Vaupel, 1991). The subunit composition of GABA (A) receptor differs in rat uteri throughout gestation and just before labour in humans, a decline in GABA receptor transcripts is observed (Bethin *et al.*, 2003, Charpigny *et al.*, 2003). TRH-R is expressed in term myometrium. (Fukusumi *et al.*, 1995) have previously reported high level of TRH-R mRNA in uterus but whether TRH plays an important role in the female reproductive tract remains to be elucidated. PGs, in conjunction with their numerous receptors activate multiple signaling pathways and exert multifaceted actions in myometrium (Hertelendy & Zakar, 2004). Thus, EP3 transcripts are down-regulated in the term pregnant myometrium (Bethin *et al.*, 2003, Charpigny *et al.*, 2003). However, the significance of these changes remains to be established, because of the known existence of EP3 splice variants.

Future strategies for integrative analysis of myometrial functions

A question is what could be the composition of an “ideal” genomic array for the study of parturition? The answer is not unique. Undedicated or "generic" microarrays seem heuristically more potent to explore and detect new interesting metabolic pathways than specialized arrays could do (Tierney *et al.*, 2003). Inversely, a myometrium-dedicated microarray is unlikely to provide essential clues on the mechanisms of parturition for the following reasons.

Gene module analysis, as exemplified in this review, searches for coordinate regulation of genes that belong to a priori defined gene modules. A statistical test performed for each module relative to all other genes on the microarray calculates whether the degree of coordinate regulation is more than one would expect by chance. Therefore, a module of genes involved in a physiological process may be significantly down-regulated whereas each gene in the module under study may be transcriptionally down regulated by say only 20%, and thus not clearly detected at the individual gene level (Tierney *et al.*, 2003, Wong & Chang, 2005). In addition, beside classical ideas regarding trans-regulation of gene expression, a greater number of hypotheses generated from regulatory networks analysis or cis-regulatory DNA elements analysis can be validated today in a high throughput fashion using chromatin-immunoprecipitation followed by microarray analysis (ChIP-chip). For example, a large fraction of genes transcribed in the liver and pancreas have been found to bind HNF4, providing a molecular explanation for the role of HNF4 mutations and polymorphisms in hereditary and sporadic forms of diabetes mellitus (Odom *et al.*, 2004).

In majority, transcripts are not tissue-restricted, but are present to varying degrees in a wide variety of cell types although there are exceptions, like myosin heavy chain which is primarily found in smooth muscle cells. As such, it is not necessary to create arrays only from cDNAs

obtained from dedicated libraries. Many sources of clones can be used for array analyses including microarrays purchased from companies (e. g. Incyte, Affymetrix, Clontech, etc.), which consist of cDNA clones or oligonucleotides that cover a large percentage of the transcripts (known and ESTs) present in public databases (Juhasz *et al.*, 2002). Only in the context of a well defined cell population, DNA microarray data can be used in a comprehensive analysis aimed at identifying the shared and unique molecular 'modules' underlying a pathological process. Thus, in peripheral organs, laser microdissection of their several constitutive compartments may reveal distinct repertoires of apoptosis-associated genes, chemokines and chemokine receptors in these compartments (Shen *et al.*, 2004).

The microarray technology made possible to widen our comprehension of the major structural and metabolic transformations which affect the myometrium from the very beginning of pregnancy until parturition. The results have proven to be coherent with the mass of information which was accumulated previously, primarily on the basis of the study of selected critical factors. Although still limited in number, the recent studies highlight the activation of precise signaling pathways, some of which may had been under evaluated. Thus, the remodelling and maturation processes that the uterus undergoes in pregnancy appear clearly as phenomenon's which last during the full course of gestation. This is attested by the nature of best represented signaling pathways, in comparisons of the non pregnant-uterus versus term uterus and the comparisons of the preterm uterus versus the term uterus in labor. Comparatively, the onset of labor is perhaps the phenomenon which remains the least well described by these methods of analysis, possibly because it is a phenomenon occurring in a too short window to have been grasped by the few studies carried out up to now. Whatever it may be, genomic studies constitute a necessary first step of orientation which should lead to a more elaborate hierarchical vision of the physiological mechanisms of gestation, in particular

by establishing new links between the generic signaling pathways that are activated during the normal or pathological gestation. Genomic studies also represent an indicative step that will need to be correlated with a systematic proteomic analysis of the myometrium. The latter will undoubtedly develop in a very near future.

Building comprehensive strategies of genomic and proteomic analysis to explore physiologic functions remains today a challenge to which very few research groups have devoted their energy, and so far only for biological functions unrelated to the pregnant uterus. The examples reported today indicate that this will be hardly accessible without the integrated cooperation of several groups (Ho *et al.*, 2003).

References

- Adolfsson PI, Dahle LO, Berg G & Svensson SP 1998 Characterization of alpha2-adrenoceptor subtypes in pregnant human myometrium. *Gynecol Obstet Invest* **45** 145-150.
- Aguan K, Carvajal JA, Thompson LP & Weiner CP 2000 Application of a functional genomics approach to identify differentially expressed genes in human myometrium during pregnancy and labour. *Mol Hum Reprod* **6** 1141-1145.
- Arntzen KJ, Kjollesdal AM, Halgunset J, Vatten L & Austgulen R 1998 TNF, IL-1, IL-6, IL-8 and soluble TNF receptors in relation to chorioamnionitis and premature labor. *J Perinat Med* **26** 17-26.
- Bailey J & Europe-Finner GN 2005 Identification of human myometrial target genes of the c-Jun NH2-terminal kinase (JNK) pathway: the role of activating transcription factor 2 (ATF2) and a novel spliced isoform ATF2-small. *J Mol Endocrinol* **34** 19-35.

- 520 Bartlett SR, Sawdy R & Mann GE 1999 Induction of cyclooxygenase-2 expression in human
 521 myometrial smooth muscle cells by interleukin-1beta: involvement of p38 mitogen-activated
 522 protein kinase. *J Physiol* **520 Pt 2** 399-406.
- 523 Bethin KE, Nagai Y, Sladek R, Asada M, Sadovsky Y, Hudson TJ & Muglia LJ 2003
 524 Microarray analysis of uterine gene expression in mouse and human pregnancy. *Mol*
 525 *Endocrinol* **17** 1454-1469.
- 526 Chan EC, Fraser S, Yin S, Yeo G, Kwek K, Fairclough RJ & Smith R 2002 Human myometrial
 527 genes are differentially expressed in labor: a suppression subtractive hybridization study. *J*
 528 *Clin Endocrinol Metab* **87** 2435-2441.
- 529 Charpigny G, Leroy MJ, Breuiller-Fouche M, Tanfin Z, Mhaouty-Kodja S, Robin P, Leiber D,
 530 Cohen-Tannoudji J, Cabrol D, Barberis C *et al.* 2003 A functional genomic study to identify
 531 differential gene expression in the preterm and term human myometrium. *Biol Reprod* **68**
 532 2289-2296.
- 533 Chegini N, Tang XM & Ma C 1999 Regulation of transforming growth factor-beta1 expression
 534 by granulocyte macrophage-colony-stimulating factor in leiomyoma and myometrial smooth
 535 muscle cells. *J Clin Endocrinol Metab* **84** 4138-4143.
- 536 Chen HW, Chen JJ, Tzeng CR, Li HN, Chang SJ, Cheng YF, Chang CW, Wang RS, Yang PC &
 537 Lee YT 2002 Global analysis of differentially expressed genes in early gestational decidua
 538 and chorionic villi using a 9600 human cDNA microarray. *Mol Hum Reprod* **8** 475-484.
- 539 Claverie JM 1999 Computational methods for the identification of differential and coordinated
 540 gene expression. *Hum Mol Genet* **8** 1821-1832.
- 541 Dallot E, Mehats C, Oger S, Leroy MJ & Breuiller-Fouche M 2005 A role for PKCzeta in the
 542 LPS-induced translocation NF-kappaB p65 subunit in cultured myometrial cells. *Biochimie*
 543 **87** 513-521.

- 544 Di Liberto G, Dallot E, Eude-Le Parco I, Cabrol D, Ferre F & Breuiller-Fouche M 2003 A
 545 critical role for PKC zeta in endothelin-1-induced uterine contractions at the end of
 546 pregnancy. *Am J Physiol Cell Physiol* **285** C599-607.
- 547 Douglas AJ, Clarke EW & Goldspink DF 1988 Influence of mechanical stretch on growth and
 548 protein turnover of rat uterus. *Am J Physiol* **254** E543-548.
- 549 Draghici S, Khatri P, Martins RP, Ostermeier GC & Krawetz SA 2003 Global functional
 550 profiling of gene expression. *Genomics* **81** 98-104.
- 551 Dunckley T, Coon KD & Stephan DA 2005 Discovery and development of biomarkers of
 552 neurological disease. *Drug Discov Today* **10** 326-334.
- 553 Esplin MS, Fausett MB, Peltier MR, Hamblin S, Silver RM, Branch DW, Adashi EY & Whiting
 554 D 2005 The use of cDNA microarray to identify differentially expressed labor-associated
 555 genes within the human myometrium during labor. *Am J Obstet Gynecol* **193** 404-413.
- 556 Eude I, Dallot E, Ferre F & Breuiller-Fouche M 2002 Protein kinase Calpha is required for
 557 endothelin-1-induced proliferation of human myometrial cells. *Biol Reprod* **66** 44-49.
- 558 Fukuhara K, Kariya M, Kita M, Shime H, Kanamori T, Kosaka C, Orii A, Fujita J & Fujii S
 559 2002 Secreted frizzled related protein 1 is overexpressed in uterine leiomyomas, associated
 560 with a high estrogenic environment and unrelated to proliferative activity. *J Clin Endocrinol*
 561 *Metab* **87** 1729-1736.
- 562 Fukusumi S, Ogi K, Onda H & Hinuma S 1995 Distribution of thyrotropin-releasing hormone
 563 receptor mRNA in rat peripheral tissues. *Regul Pept* **57** 115-121.
- 564 Ganesh SK, Skelding KA, Mehta L, O'Neill K, Joo J, Zheng G, Goldstein J, Simari R, Billings
 565 E, Geller NL *et al.* 2004 Rationale and study design of the CardioGene Study: genomics of in-
 566 stent restenosis. *Pharmacogenomics* **5** 952-1004.

- 567 Gargiulo AR, Khan-Dawood FS & Dawood MY 1997 Epidermal growth factor receptors in
 568 uteroplacental tissues in term pregnancy before and after the onset of labor. *J Clin Endocrinol*
 569 *Metab* **82** 113-117.
- 570 Girotti M & Zingg HH 2003 Gene expression profiling of rat uterus at different stages of
 571 parturition. *Endocrinology* **144** 2254-2265.
- 572 Gogarten W, Emala CW, Lindeman KS & Hirshman CA 2001 Oxytocin and lysophosphatidic
 573 acid induce stress fiber formation in human myometrial cells via a pathway involving Rho-
 574 kinase. *Biol Reprod* **65** 401-406.
- 575 Grammatopoulos DK, Randeva HS, Levine MA, Katsanou ES & Hillhouse EW 2000 Urocortin,
 576 but not corticotropin-releasing hormone (CRH), activates the mitogen-activated protein kinase
 577 signal transduction pathway in human pregnant myometrium: an effect mediated via R1alpha
 578 and R2beta CRH receptor subtypes and stimulation of Gq-proteins. *Mol Endocrinol* **14** 2076-
 579 2091.
- 580 Guha M & Mackman N 2001 LPS induction of gene expression in human monocytes. *Cell*
 581 *Signal* **13** 85-94.
- 582 Havelock JC, Keller P, Muleba N, Mayhew BA, Casey BM, Rainey WE & Word RA 2005
 583 Human myometrial gene expression before and during parturition. *Biol Reprod* **72** 707-719.
- 584 Heinrich PC, Behrmann I, Muller-Newen G, Schaper F & Graeve L 1998 Interleukin-6-type
 585 cytokine signalling through the gp130/Jak/STAT pathway. *Biochem J* **334** (Pt 2) 297-314.
- 586 Hertelendy F & Zakar T 2004 Regulation of myometrial smooth muscle functions. *Curr Pharm*
 587 *Des* **10** 2499-2517.
- 588 Hertelendy F, Molnar M & Romero R 2002 Interferon gamma antagonizes interleukin-1beta-
 589 induced cyclooxygenase-2 expression and prostaglandin E(2) production in human
 590 myometrial cells. *J Soc Gynecol Investig* **9** 215-219.

- 591 Hewitt SC, Collins J, Grissom S, Deroo B & Korach KS 2005 Global uterine genomics in vivo:
 592 microarray evaluation of the estrogen receptor alpha-growth factor cross-talk mechanism. *Mol*
 593 *Endocrinol* **19** 657-668.
- 594 Ho M, Yang E, Matcuk G, Deng D, Sampas N, Tsalenko A, Tabibiazar R, Zhang Y, Chen M,
 595 Talbi S *et al.* 2003 Identification of endothelial cell genes by combined database mining and
 596 microarray analysis. *Physiol Genomics* **13** 249-262.
- 597 Juhasz O, Zhu Y, Garg R, Anisimov SV & Boheler KR 2002 Analysis of altered genomic
 598 expression profiles in the senescent and diseased myocardium using cDNA microarrays. *Eur*
 599 *J Heart Fail* **4** 687-697.
- 600 Kanehisa M 1997 A database for post-genome analysis. *Trends Genet* **13** 375-376.
- 601 Kanehisa M & Goto S 2000 KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*
 602 *Res* **28** 27-30.
- 603 Kato HD, Terao Y, Ogawa M, Matsuda T, Arima T, Kato K, Yong Z & Wake N 2002 Growth-
 604 associated gene expression profiles by microarray analysis of trophoblast of molar
 605 pregnancies and normal villi. *Int J Gynecol Pathol* **21** 255-260.
- 606 Kayisli UA, Mahutte NG & Arici A 2002 Uterine chemokines in reproductive physiology and
 607 pathology. *Am J Reprod Immunol* **47** 213-221.
- 608 Keelan JA, Blumenstein M, Helliwell RJ, Sato TA, Marvin KW & Mitchell MD 2003 Cytokines,
 609 prostaglandins and parturition--a review. *Placenta* **24 Suppl A** S33-46.
- 610 Khatri P & Draghici S 2005 Ontological analysis of gene expression data: current tools,
 611 limitations, and open problems. *Bioinformatics* **21** 3587-3595.
- 612 Khatri P, Sellamuthu S, Malhotra P, Amin K, Done A & Draghici S 2005 Recent additions and
 613 improvements to the Onto-Tools. *Nucleic Acids Res* **33** W762-765.

- 614 Kimura A, Ohmichi M, Takeda T, Kurachi H, Ikegami H, Koike K, Masuhara K, Hayakawa J,
 615 Kanzaki T, Kobayashi M *et al.* 1999 Mitogen-activated protein kinase cascade is involved in
 616 endothelin-1-induced rat puerperal uterine contraction. *Endocrinology* **140** 722-731.
- 617 Klotz DM, Hewitt SC, Ciana P, Raviscioni M, Lindzey JK, Foley J, Maggi A, DiAugustine RP
 618 & Korach KS 2002 Requirement of estrogen receptor-alpha in insulin-like growth factor-1
 619 (IGF-1)-induced uterine responses and in vivo evidence for IGF-1/estrogen receptor cross-
 620 talk. *J Biol Chem* **277** 8531-8537.
- 621 Leppert PC 1998 Proliferation and apoptosis of fibroblasts and smooth muscle cells in rat uterine
 622 cervix throughout gestation and the effect of the antiprogesterone onapristone. *Am J Obstet*
 623 *Gynecol* **178** 713-725.
- 624 Leroy MJ, Lugnier C, Merezak J, Tanguy G, Olivier S, Le Bec A & Ferre F 1994 Isolation and
 625 characterization of the rolipram-sensitive cyclic AMP-specific phosphodiesterase (type IV
 626 PDE) in human term myometrium. *Cell Signal* **6** 405-412.
- 627 Leroy MJ, Mehats C, Duc-Goiran P, Tanguy G, Robert B, Dallot E, Mignot TM, Grange G &
 628 Ferre F 1999 Effect of pregnancy on PDE4 cAMP-specific phosphodiesterase messenger
 629 ribonucleic acid expression in human myometrium. *Cell Signal* **11** 31-37.
- 630 Lewis SE 2005 Gene Ontology: looking backwards and forwards. *Genome Biol* **6** 103.
- 631 Li Y, Je HD, Malek S & Morgan KG 2004 Role of ERK1/2 in uterine contractility and preterm
 632 labor in rats. *Am J Physiol Regul Integr Comp Physiol* **287** R328-335.
- 633 Liu Y, Dong W, Chen L, Xiang R, Xiao H, De G, Wang Z & Qi Y 2004 BCL10 mediates
 634 lipopolysaccharide/toll-like receptor-4 signaling through interaction with Pellino2. *J Biol*
 635 *Chem* **279** 37436-37444.
- 636 Loch-Caruso RK, Criswell KA, Grindatti CM & Brant KA 2003 Sustained inhibition of rat
 637 myometrial gap junctions and contractions by lindane. *Reprod Biol Endocrinol* **1** 62.

- 638 Lopez Bernal A & TambyRaja RL 2000 Preterm labour. *Baillieres Best Pract Res Clin Obstet*
639 *Gynaecol* **14** 133-153.
- 640 Macphee DJ & Lye SJ 2000 Focal adhesion signaling in the rat myometrium is abruptly
641 terminated with the onset of labor. *Endocrinology* **141** 274-283.
- 642 Majewska MD & Vaupel DB 1991 Steroid control of uterine motility via gamma-aminobutyric
643 acidA receptors in the rabbit: a novel mechanism? *J Endocrinol* **131** 427-434.
- 644 Mangioni S, Vigano P, Lattuada D, Abbiati A, Vignali M & Di Blasio AM 2005 Overexpression
645 of the Wnt5b gene in leiomyoma cells: implications for a role of the Wnt signaling pathway in
646 the uterine benign tumor. *J Clin Endocrinol Metab* **90** 5349-5355.
- 647 Marvin KW, Keelan JA, Eykholt RL, Sato TA & Mitchell MD 2002a Expression of angiogenic
648 and neurotrophic factors in the human amnion and choriodecidua. *Am J Obstet Gynecol* **187**
649 728-734.
- 650 Marvin KW, Keelan JA, Eykholt RL, Sato TA & Mitchell MD 2002b Use of cDNA arrays to
651 generate differential expression profiles for inflammatory genes in human gestational
652 membranes delivered at term and preterm. *Mol Hum Reprod* **8** 399-408.
- 653 Mehats C, Tanguy G, Dallot E, Cabrol D, Ferre F & Leroy MJ 2001 Is up-regulation of
654 phosphodiesterase 4 activity by PGE2 involved in the desensitization of beta-mimetics in late
655 pregnancy human myometrium? *J Clin Endocrinol Metab* **86** 5358-5365.
- 656 Mehats C, Tanguy G, Paris B, Robert B, Pernin N, Ferre F & Leroy MJ 2000 Pregnancy induces
657 a modulation of the cAMP phosphodiesterase 4-conformers ratio in human myometrium:
658 consequences for the utero-relaxant effect of PDE4-selective inhibitors. *J Pharmacol Exp*
659 *Ther* **292** 817-823.
- 660 Mhaouty-Kodja S, Houdeau E & Legrand C 2004 Regulation of myometrial phospholipase C
661 system and uterine contraction by beta-adrenergic receptors in midpregnant rat. *Biol Reprod*
662 **70** 570-576.

- 663 Moore F, Da Silva C, Wilde JI, Smarason A, Watson SP & Lopez Bernal A 2000 Up-regulation
 664 of p21- and RhoA-activated protein kinases in human pregnant myometrium. *Biochem*
 665 *Biophys Res Commun* **269** 322-326.
- 666 Morimoto T, Head JR, MacDonald PC & Casey ML 1998 Thrombospondin-1 expression in
 667 human myometrium before and during pregnancy, before and during labor, and in human
 668 myometrial cells in culture. *Biol Reprod* **59** 862-870.
- 669 Muhle RA, Pavlidis P, Grundy WN & Hirsch E 2001 A high-throughput study of gene
 670 expression in preterm labor with a subtractive microarray approach. *Am J Obstet Gynecol* **185**
 671 716-724.
- 672 Nishimura M & Naito S 2005 Tissue-specific mRNA expression profiles of human toll-like
 673 receptors and related genes. *Biol Pharm Bull* **28** 886-892.
- 674 Odom DT, Zizlsperger N, Gordon DB, Bell GW, Rinaldi NJ, Murray HL, Volkert TL, Schreiber
 675 J, Rolfe PA, Gifford DK *et al.* 2004 Control of pancreas and liver gene expression by HNF
 676 transcription factors. *Science* **303** 1378-1381.
- 677 Ohmichi M, Koike K, Kimura A, Masuhara K, Ikegami H, Ikebuchi Y, Kanzaki T, Touhara K,
 678 Sakaue M, Kobayashi Y *et al.* 1997 Role of mitogen-activated protein kinase pathway in
 679 prostaglandin F₂α-induced rat puerperal uterine contraction. *Endocrinology* **138** 3103-
 680 3111.
- 681 Oldenhof AD, Shynlova OP, Liu M, Langille BL & Lye SJ 2002 Mitogen-activated protein
 682 kinases mediate stretch-induced c-fos mRNA expression in myometrial smooth muscle cells.
 683 *Am J Physiol Cell Physiol* **283** C1530-1539.
- 684 Osman I, Young A, Ledingham MA, Thomson AJ, Jordan F, Greer IA & Norman JE 2003
 685 Leukocyte density and pro-inflammatory cytokine expression in human fetal membranes,
 686 decidua, cervix and myometrium before and during labour at term. *Mol Hum Reprod* **9** 41-45.

- 687 Otun HA, MacDougall MW, Bailey J, Europe-Finner GN & Robson SC 2005 Spatial and
 688 temporal expression of the myometrial mitogen-activated protein kinases p38 and ERK1/2 in
 689 the human uterus during pregnancy and labor. *J Soc Gynecol Investig* **12** 185-190.
- 690 Ozaki H, Yasuda K, Kim YS, Egawa M, Kanzaki H, Nakazawa H, Hori M, Seto M & Karaki H
 691 2003 Possible role of the protein kinase C/CPI-17 pathway in the augmented contraction of
 692 human myometrium after gestation. *Br J Pharmacol* **140** 1303-1312.
- 693 Papadopoulou N, Chen J, Randeva HS, Levine MA, Hillhouse EW & Grammatopoulos DK 2004
 694 Protein kinase A-induced negative regulation of the corticotropin-releasing hormone R1alpha
 695 receptor-extracellularly regulated kinase signal transduction pathway: the critical role of
 696 Ser301 for signaling switch and selectivity. *Mol Endocrinol* **18** 624-639.
- 697 Phaneuf S, Carrasco MP, Europe-Finner GN, Hamilton CH & Lopez Bernal A 1996 Multiple G
 698 proteins and phospholipase C isoforms in human myometrial cells: implication for oxytocin
 699 action. *J Clin Endocrinol Metab* **81** 2098-2103.
- 700 Qi M & Elion EA 2005 Formin-induced actin cables are required for polarized recruitment of the
 701 Ste5 scaffold and high level activation of MAPK Fus3. *J Cell Sci* **118** 2837-2848.
- 702 Rehman KS, Yin S, Mayhew BA, Word RA & Rainey WE 2003 Human myometrial adaptation
 703 to pregnancy: cDNA microarray gene expression profiling of myometrium from non-pregnant
 704 and pregnant women. *Mol Hum Reprod* **9** 681-700.
- 705 Robin P, Boulven I, Desmyter C, Harbon S & Leiber D 2002 ET-1 stimulates ERK signaling
 706 pathway through sequential activation of PKC and Src in rat myometrial cells. *Am J Physiol*
 707 *Cell Physiol* **283** C251-260.
- 708 Robin P, Boulven I, Bole-Feysot C, Tanfin Z & Leiber D 2004 Contribution of PKC-dependent
 709 and -independent processes in temporal ERK regulation by ET-1, PDGF, and EGF in rat
 710 myometrial cells. *Am J Physiol Cell Physiol* **286** C798-806.

- 711 Romero R, Kuivaniemi H & Tromp G 2002 Functional genomics and proteomics in term and
 712 preterm parturition. *J Clin Endocrinol Metab* **87** 2431-2434.
- 713 Ruzycky AL 1998 Down-regulation of the mitogen-activated protein kinase cascade
 714 immediately before parturition in the rat myometrium. *J Soc Gynecol Investig* **5** 304-310.
- 715 Salomonis N, Cotte N, Zambon AC, Pollard KS, Vranizan K, Doniger SW, Dolganov G &
 716 Conklin BR 2005 Identifying genetic networks underlying myometrial transition to labor.
 717 *Genome Biol* **6** R12.
- 718 Sehringer B, Zahradnik HP, Simon M, Ziegler R, Noethling C & Schaefer WR 2004 mRNA
 719 expression profiles for corticotrophin-releasing hormone, urocortin, CRH-binding protein and
 720 CRH receptors in human term gestational tissues determined by real-time quantitative RT-
 721 PCR. *J Mol Endocrinol* **32** 339-348.
- 722 Shen Y, Iqbal J, Xiao L, Lynch RC, Rosenwald A, Staudt LM, Sherman S, Dybkaer K, Zhou G,
 723 Eudy JD *et al.* 2004 Distinct gene expression profiles in different B-cell compartments in
 724 human peripheral lymphoid organs. *BMC Immunol* **5** 20.
- 725 Shynlova O, Tsui P, Dorogin A, Chow M & Lye SJ 2005 Expression and localization of alpha-
 726 smooth muscle and gamma-actins in the pregnant rat myometrium. *Biol Reprod* **73** 773-780.
- 727 Sooranna SR, Lee Y, Kim LU, Mohan AR, Bennett PR & Johnson MR 2004 Mechanical stretch
 728 activates type 2 cyclooxygenase via activator protein-1 transcription factor in human
 729 myometrial cells. *Mol Hum Reprod* **10** 109-113.
- 730 Stein RC & Waterfield MD 2000 PI3-kinase inhibition: a target for drug development? *Mol Med*
 731 *Today* **6** 347-357.
- 732 Tashima LS, Yamamoto SY, Yasuda M, Millar LK & Bryant-Greenwood GD 2002 Decidual
 733 relaxins: gene and protein up-regulation in preterm premature rupture of the membranes by
 734 complementary DNA arrays and quantitative immunocytochemistry. *Am J Obstet Gynecol*
 735 **187** 785-797.

- 736 Tierney EP, Tulac S, Huang ST & Giudice LC 2003 Activation of the protein kinase A pathway
 737 in human endometrial stromal cells reveals sequential categorical gene regulation. *Physiol*
 738 *Genomics* **16** 47-66.
- 739 Wang H & Hirsch E 2003 Bacterially-induced preterm labor and regulation of prostaglandin-
 740 metabolizing enzyme expression in mice: the role of toll-like receptor 4. *Biol Reprod* **69**
 741 1957-1963.
- 742 Wang Y, Matsuo H, Kurachi O & Maruo T 2002 Down-regulation of proliferation and up-
 743 regulation of apoptosis by gonadotropin-releasing hormone agonist in cultured uterine
 744 leiomyoma cells. *Eur J Endocrinol* **146** 447-456.
- 745 Williams SJ, White BG & MacPhee DJ 2005 Expression of alpha5 integrin (Itga5) is elevated in
 746 the rat myometrium during late pregnancy and labor: implications for development of a
 747 mechanical syncytium. *Biol Reprod* **72** 1114-1124.
- 748 Wong DJ & Chang HY 2005 Learning more from microarrays: insights from modules and
 749 networks. *J Invest Dermatol* **125** 175-182.
- 750 Wu WX, Zhang Q, Ma XH, Unno N & Nathanielsz PW 1999 Suppression subtractive
 751 hybridization identified a marked increase in thrombospondin-1 associated with parturition in
 752 pregnant sheep myometrium. *Endocrinology* **140** 2364-2371.
- 753 Yu JT & Lopez Bernal A 1998 The cytoskeleton of human myometrial cells. *J Reprod Fertil* **112**
 754 185-198.
- 755 Zhong M, Yang M & Sanborn BM 2003 Extracellular signal-regulated kinase 1/2 activation by
 756 myometrial oxytocin receptor involves Galpha(q)Gbetagamma and epidermal growth factor
 757 receptor tyrosine kinase activation. *Endocrinology* **144** 2947-2956.
- 758 Zhong M, Ku CY & Sanborn BM 2005 Pathways used by relaxin to regulate myometrial
 759 phospholipase C. *Ann N Y Acad Sci* **1041** 300-304.
- 760

Table 1

Actin cytoskeleton			Focal adhesion			Adherens junctions			Tight junctions		
NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³
15	6	0	18	6	3	11	3	0	10	0	0
ACTN4(+)	EGF(-)		ACTN4(+)	INSR(+)	ILK(+)	ACTN4(+)	INSR(+)		ACTN4(+)		
CD14(+)	ITGAV(-)		ARHGAP5(+)	ITGB3(-)	THBS1(+)	CTNNA1(+)	PTPRB5(-)		CTNNA1(+)		
EGFR(+)	ITGB3(-)		CAV2(+)	ITGB6(-)	TNC(+)	EGFR(+)	PTPN1(-)		GNAI2(+)		
FGFR1(+)	ITGB6(-)		COL1A2(+)	MAPK10(-)		FGFR1(+)			HCLS1(+)		
HEM1(+)	MATK(-)		EGFR(+)	PIK3CB(-)		NLK(+)			PRKCB1(+)		
ITGAM(+)	PIK3CB(-)		FLNA(+)	RAP1A(-)		PTPN6(+)			PRKCD(+)		
ITGAV(+)			ITGB5(+)			PTPRJ(+)			PRKCG(+)		
ITGB5(+)			PIK3CA(+)			RAC2(+)			RAB13(+)		
PFN1(+)			PIP5K1C(+)			SMAD2(+)			RRAS(+)		
PIK3CA(+)			PRKCB1(+)			TCF7L2(+)			SPTAN1(+)		
PIP5K1C(+)			PRKCG(+)			VCL(+)					
RAC2(+)			ROCK1(+)								
ROCK1(+)			RRAS(+)								
RRAS(+)			TNC(+)								
VCL(+)			VASP(+)								
			VCL(+)								
			VWF(+)								
			ZYX(+)								

Table 1 legend

The main KEGG (<http://www.genome.jp/kegg/>) metabolic pathways, regulatory pathways or molecular complexes that were found to be the best represented in the myometrium on the basis of genes detected in genomic studies according to ¹NP-TNIL (Rehman *et al.*, 2003), to ²PT-TNIL (Bethin *et al.*, 2003, Charpigny *et al.*, 2003) or to ³TNIL-TIL (Esplin *et al.*, 2005, Havelock *et al.*, 2005), during the cycle or pregnancy. All gene

transcripts showed at least a 2 fold or more, up- (+) or down-regulation (-) after paired comparisons of the following stages: non-pregnant (NP) versus term not in labor (TNIL), preterm (PT) versus TNIL, or TNIL versus term in labor (TIL). Number of genes represented in pathways and physiological stages, out of the total number of genes (n=118) sorted as belonging to these pathways, “0” means no gene represented for the corresponding pathway. OMIM nomenclature adopted for gene names (<http://www.ncbi.nlm.nih.gov/omim/>).

Table 2

Cytokine-cytokine receptors			Toll-like receptors			TGF-beta signaling pathway			Wnt signaling pathway		
NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³
8	12	5	5	5	1	6	2	3	7	2	1
CCL19(+)	BLR1(+)	CCL13(+)	CD14(+)		CXCL10(+)	ACVRL1(+)	THBS2(+)	TGFB2(+)	NLK(+)	MAPK10(-)	VANGL1(+)
CCL21(+)	CCL13(+)	CCL18(+)	IL6(+)			BMP6(+)	TNF(-)	THBS1(+)	PRKCB1(+)	TP53(+)	
CSF1R(+)	CXCR4(+)	CXCL10(+)	PIK3CA(+)		MAP2K4(-)	ROCK1(+)		THBS2(+)	PRKCG(+)		
EGFR(+)	IL9(+)	TGFB2(+)	RAC2(+)		MAPK10(-)	SMAD2(+)			RAC2(+)		
IL15RA(+)	TNFRSF6(+)	TNFRSF11B(+)	STAT1(+)		PIK3CB(-)	TFDP1(+)			ROCK1(+)		
IL6(+)	CLC(-)				TNF(-)	THBS2(+)			SMAD2(+)		
IL6ST(+)	EGF(-)								TCF7L2(+)		
TNFRSF11B(+)	IFNGR1(-)										
	IL6ST(-)										
	LIFR(-)										
	TNF(-)										
	TNFSF4(-)										

Table 2 legend

See Table 1 legend

Table 3

MAPK signaling pathway			Phosphatidylinositol signaling pathway			JAK-STAT signaling pathway		
NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³
15	12	4	5	8	0	6	7	1
CD14(+)	EGF(+)	BDNF(+)	IMPA2(+)	CALM1(-)		IL15RA(+)	IL9(+)	IL13(+)
EGFR(+)	IKBKB(+)	DUSP5(+)	PIK3CA(+)	CALM3(-)		IL6(+)	PIM1(+)	
FGFR1(+)	MAP2K4(+)	NR4A1(+)	PIP5K1C(+)	DGKZ(-)		IL6ST(+)	CLC(-)	
FLNA(+)	MAPK10(+)	TGFB2(+)	PRKCB1(+)	ITPR1(-)		PIK3CA(+)	IFNGR1(-)	
HSPA5(+)	PLA2G5(+)		PRKCG(+)	ITPR1(-)		PTPN6(+)	IL6ST(-)	
HSPA8(+)	RAP1A(+)			PIK3CB(-)		STAT1(+)	LIFR(-)	
MAP3K5(+)	RASA1(+)			PLCG1(-)			PIK3CB(-)	
NLK(+)	TNF(+)			PLCG2(-)				
PLA2G2A(+)	TP53(+)							
PRKCB1(+)	PLA2G2A(-)							
PRKCG(+)	PPP5C(-)							
RAC2(+)	TNFRSF6(-)							
RASGRP2(+)								
RRAS(+)								
YWHAZ(+)								

Table 3 legend

See Table 1 legend

Table 4

Apoptosis		Calcium signaling pathway			Neuroactive ligand-receptors		
NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ²	TNIL-TIL ³	NP-TNIL ¹	PT-TNIL ² TNIL-TIL ³
0	6	0	8	8	0	6	6 0
	TNFRSF6(+)		EGFR(+)	BST1(+)		ADORA3(+)	GLRB(+)
	TP53(+)		GNAQ(+)	CALM1(-)		ADRA2C(+)	GABRB1(-)
	DFFA(-)		LYN(+)	CALM3(-)		C3AR1(+)	GABRG2(-)
	IRAK1(-)		NFKBIB(+)	ITPR1(-)		EDN1(+)	PTGER3(-)
	PIK3CB(-)		PHKA1(+)	PLCG1(-)		GRM2(+)	TACR2(-)
	TNF(-)		PRKCB1(+)	PLCG2(-)		NPY(+)	TRHR(-)
			PRKCG(+)	PTGER3(-)			
			PYGL(+)	TACR2(-)			

Table 4 legend

See Table 1 legend

Figure legends

Figure 1

Main KEGG pathways (<http://www.genome.jp/kegg/>) represented in the myometrium on the basis of the genes detected in genomic studies. Gene transcripts pertaining to these pathways were either up-regulated (red) or down-regulated (green) by a two fold or more factor. White boxes denote insufficiently documented links in context. See Tables 1-4 for details. (A) Comparison between the non-pregnant (NP) and term not in labor (TNIL) uterus (Rehman et al., 2003). (B) Comparison between the preterm (PT) uterus and term in labor uterus (TNIL) (Bethin et al., 2003, Charpigny et al., 2003) or between TNIL and term in labor uterus (TIL) (Esplin et al., 2005, Havelock et al., 2005). Five pathways (Focal adhesion, MAPK, TGF-beta, JAK-STAT and apoptosis) are shown with their known links to membrane receptors (Wnt, TGF-beta, cytokines and Toll-like receptors). Three additional pathways (actin cytoskeleton, phosphatidylinositol and calcium signaling) are over-expressed in gestation by comparison to the non-pregnant state. THBS1, thrombospondin-1 (inhibitory).

Figure 2

Main KEGG pathways (<http://www.genome.jp/kegg/>) represented in the myometrium on the basis of the genes detected in genomic studies. Gene transcripts pertaining to these pathways were either up-regulated (red) or down-regulated (green) by a two fold or more factor. White boxes denote insufficiently documented links in context. See Tables 1-4 for details. (A) Comparison between the non-pregnant (NP) and term not in labour (TNIL) uterus (Rehman et al., 2003). (B) Comparison between the preterm (PT) uterus and term in labor uterus

26 (TNIL) (Bethin et al., 2003, Charpigny et al., 2003) or between TNIL and term in labor uterus
27 (TIL) (Esplin et al., 2005, Havelock et al., 2005). Ten pathways (focal adhesion, adherens
28 junctions, tight junctions, actin cytoskeleton, phosphatidylinositol and calcium signaling,
29 MAPK, TGF-beta, JAK-STAT and apoptosis) are shown with their known links to cytokine,
30 growth factors (GF) and G-protein coupled membrane receptors. CAMs, cell adhesion
31 molecules, ECM, extra-cellular matrix.

32

Figure 1




Figure 2

